



รายงานการวิจัยฉบับสมบูรณ์

Isolation of Endophytic Fungi from some Orchid Varieties



E077877

ผ.ศ. ดร.ศรายุทธ ผลโพธิ์

นายวรรณะ ด้วง

สาขา.....
เลขทะเบียน 077877
วัน เดือน ปี 22 ก.ย. 2559

12804708
.b.....
.i.....

ได้รับทุนสนับสนุนการวิจัยจากงบประมาณปี ประจำปีงบประมาณ 2557

คณะเทคโนโลยีการเกษตร

สถาบันเทคโนโลยีพระจอมเกล้าเจ้าคุณทหารลาดกระบัง

This material is reserved for educational use only, not allowed for commercial use.

Forbidden to modify the content, and cite the document when use.

Isolation of Endophytic Fungi from some Orchid Varieties

ชื่อโครงการ Isolation of Endophytic Fungi from some Orchid Varieties

แหล่งเงิน เงินงบประมาณ คณะเทคโนโลยีการเกษตร สถาบันเทคโนโลยีพระจอมเกล้าเจ้าคุณทหาร

ลาดกระบัง

ประจำปีงบประมาณ 2557 จำนวนเงินที่ได้รับการสนับสนุน 100,000 บาท

ระยะเวลาทำการวิจัย 1 ปี ตั้งแต่ 1 ตุลาคม ถึง 30 กันยายน 2555

ชื่อ-สกุล หัวหน้าโครงการ ผศ.ดร.ศรายุทธ ผลโพธิ์

ผู้ร่วมโครงการวิจัย นายวรรณะ ส้ว นักศึกษาระดับปริญญาโท สาขาเกษตรศาสตร์

Abstract

First study on endophytic fungi from orchids in Chantaburi province, Thailand. Endophytic fungi were isolated from 9 orchid varieties in green house condition. Leaves and roots of *Grammatophyllum scriptum*, *Cymbidium dayanum*, *Dendrobium hercoglossum*, *Dendrobiumpalpebrae*, *Toreniafournieri*, *Doritispulcherrima*, *Dendrobium crumenatum*, *Dendrobium friedericksianum* and *Grammatophyllum specinocum*. All samples were collected then brought to laboratory. Isolation was done by surface sterilization, removed the outer surface and got inner tissues then place on water agar and transferred culture to potato dextrose agar to be pure culture. The morphological identification based on colony and agar color, liquid drops, colony shapes, the growth rate and spores characteristic. Twelve species are identified as *Chaetomium cochliodes*, *Chaetomium cupreum*, *Xylaria* sp., *Colletotrichum* sp., *Nigrospora* sp., *Phoma* sp., *Curvularia* sp., *Fusarium* sp., *Pestalotiopsis* sp., *Corynascus* sp. and *Cladosporium*-like fungi, were obtained. Most of fungi that isolated were familiar to endophytic that isolated from many plant as previous study. All of the fungi have different benefit up to using.

Keywords: Endophytic fungi, Orchids

Isolation of Endophytic Fungi from some Orchid Varieties

Acknowledgement

This research has got scholarship from King Mongkut's Institute of Technology Ladkrabang. I would like to express my sincere to thank to Faculty of Agriculture Technology, King Mongkut's Institute of Technology Ladkrabang (KMITL), Bangkok, Thailand. I also thank to the teacher and stuff of Rajamangala University Technology Tawan-ok Chantaburi campus what allowed to collect orchid samples in this research.

ผศ.ดร.ศรายุทธ ผลโพธิ์

นายวรรณะ ส้ว



This material is reserved for educational use only, not allowed for commercial use.

Forbidden to modify the content, and cite the document when use ศรายุทธ ผลโพธิ์ วรรณะ ส้ว

Isolation of Endophytic Fungi from some Orchid Varieties


Table of Content

Content	Page
Abstract	i
Acknowledgments	ii
Table of content	iii
List of figures	iv
Chapter 1 Introduction	1
Chapter 2 Review literature	3
Chapter 3 Materials and Methods	11
Chapter 4 Result and Discussion	13
Reference	20
Appendix	22



List of figures

Figures	Page
Fig. 1: <i>Chaetomium cochliodes</i> A: The colony that culture on PDA 30 days, B The spores under microscops 10X C The ascospores under microscops 40X	13
Fig. 2: <i>Chaetomium cupreum</i> A: The colony cultured on PDA 30 days B: The spores under microscops 10X and C: The ascospores under microscops 40X	13
Fig. 3: <i>Xylaria</i> sp. A: Colony culture on PDA 30 days B: Mycelia of xylaria sp. under microscops 40X	14
Fig. 4: <i>Colletotrichum</i> sp. A: The colony culture on PDA 30 days B: The spores under microscope 40X	14
Fig. 5: <i>Nigrospora</i> sp. A: The colony culture on PDA 30 days B: The hyphea and spores under microscope 40X	15
Fig. 6: <i>Phoma</i> sp. A: The colony culture on PDA 30 days and B: The spores and mycelia under microscope 40X	15
Fig. 7: <i>Curvularia</i> sp. A: The colony culture on PDA 30 days and B: The spores under microscope 40X	16
Fig. 9: <i>Pestalotiopsis</i> sp. A: The colony culture on PDA 30 days B: The conidia under microscops 40X	16
Fig. 10: <i>Corynascus</i> sp. A: The colony culture on PDA 30 days and B: The spores and ascospores under microscops 40X	16
Fig. 11: <i>Cladosporium</i> -like fungi A: The colony culture on PDA 30 days and B: The spores and mycelia under microscops 40X	17
Fig. 12: <i>Achaetomium</i> sp. A: The colony culture on PDA 30 days B: The spores under microscops 10X and C: The ascospores under microscops 40X	27



Chapter 1

Introduction

Plants are usually associated with diverse microorganisms. Endophyte microorganisms are those that colonize the healthy plant internal tissue (Stone J.K. *et al.*, 2004). The meaning of term “endophyte” is as broad as its literal definition and spectrum of potential hosts and inhabitants. Endophytes are used for both bacteria and fungi (Schulz, B. *et al.*, 2006). Modern usage of the term endophytic fungi in mycology refers to those fungi which live within leaves stems roots and other part of apparently healthy host plants and can not be seen in visually signs of infection (Stone J.K. *et al.*, 2004). Dreyfuss and Chapela (1994) estimated that there may be at least one million species of endophytic fungi alone. It means that almost all plant species are usually infected with endophytic fungi.

Recently endophytic fungi have attracted the attention from many scientists in the world as estimated that such species may useful as sources of anticancer, antidiabetic, insecticidal and immunosuppressive compounds (Strobel, G. and Daisy, B., 2003). Endophytic fungi from medicinal plants can be used for the development of drugs. For example *Fusarium oxysporum* isolated from *Catharanthus roseus* plant can produce Vinblastine and Vincristine, excellent anti-cancer drugs (Kumar A. *et al.*, 2013). Some species of endophytic fungi from Chinese medicinal plant *Actinidia macrosperma* can inhibit the tumour cells (Yin Lu *et al.*, 2011). The differences in diversity and types of endophytic fungi depend on their host plants and changes in environment (Gange *et al.*, 2007). For the beneficial role of endophyte in plants can be divided into two categories based on types of activity: growth promotion and disease control (Anu Rajan, S., 2012). It can directly enhance plant growth, increase the host plants' adaptability to such diverse conditions as dry, cold, and high-salt environments, and help to resist pathogen damage (Waller F., 2005) as well as being able to alter the community structure (Purahong W, Hyde KD., 2011) and antioxidant activity of the host plants (Hamilton CE., 2012). For example fungal endophyte, *Scolecobasidium humicola* isolated from the roots of tomato can increase the tomato biomass with the organic nitrogen sources (Rola S. Mahmoud and Kazuhiko Narisawa, 2013). They also can produce gibberellin and indoleacetic acid and promote host plant during stress condition (Waqas M. *et al.*, 2012).

Orchids are monocotyledonous plants and taxonomically belong to Order Orchidales and Family Orchidaceae. There are 15000 to 25000 of orchid species distributed all around the world, mostly can found in tropical environment. They all can divide in 2 classifications as epiphytic orchids and terrestrial orchids. In Thailand, there are about 500 genus and 800 orchid species. The special hereditary characteristic of orchid species is having many tiny seed in the pod and there are not growing and no endosperm in embryo. So to be germinated, the orchids have to depend on some fungi species to be the energy and food sources such as many nutrients that important for growth (Thamasiri K., 2005).

Orchids are among the plant groups that have aroused most widespread interest among scientists and horticulturists, both for their study and use. They have been subject to particularly high commercial demand over the past 40 years for the beauty of their structure and their vividly colored flowers, steeped in symbolism and mystery. Some countries have declared orchid species as their national flowers (Clemente M., 2009). In Thailand orchids are the most valuable product for export and make money for country. Last years (2013) the export of orchids value 563,134,221 Bart, and the country that most import Thai orchids is China, Japan, U.S.A, India and Vietnam, respectively (Source: Office of Agriculture Economics).

Fungal endophytes have now been investigated in a large number of orchid species from around the world. The traditional approach to identify the fungal endophytes of orchids has been to isolate pelotons from orchid tissues and to maintain fungal colonies in pure culture. The fungi were then identified on the basis of anatomy and morphology including such features as nucleus number, hyphal cross wall structure and spore dimensions (Perkins *et al.* 1995). Worldwide, the fungi involved with orchids are almost all members of the phylum Basidiomycota group, however many do not

This material is reserved for educational use only, not allowed for commercial use.

Forbidden to modify the content, and cite the document when use.

produce sexual spores, and are consequently assigned to the form genus *Rhizoctonia* (Rasmussen 2002). Although results may vary with the species of orchid examined, it appears that under natural conditions a particular orchid species may associate with only a few or a single fungal species. For example the *Xylaria* spp. and *Rhizotonia-like* fungi were isolate from root and leaves of genus *Lepanthes* (Bayman, P., et al., 1997). Bougoure J.J. and J.D.W. Dearnaley isolated the endophytic fungi form *Dipodium Variegatum* (Orchidaceae) found the *Russula* sp. and non-mycorrhizal Deuteromycetes.

This study aims to investigate the diversity of endophytic fungi by isolation from the roots and leave of 9 orchid varieties in Rajamangala University of Technology Tawan-ok Chantaburi campus.

State of Problem

At present the most farmers always use chemical matter such ass chemical fertilizers insecticide, cause effect to farmer's health and many toxic matter contaminate to the agriculture products. To investigate for bioactive from nature is the main project of all scientists, to replace of chemical matter. This research project is to isolate endophytic fungi from various orchid varieties, that this is the first step to find the beneficial fungi for further analysis for bio-control in orchid production.

Objectives

1. To isolate the endophytic fungi form various orchid varieties;
2. To identify the species of endophytic fungi from various orchid varieties.

Location of Research

Biocontrol research Laboratory and Mycology Section, Faculty of Agricultural Technology, King Mongkut's Institute of Technology Ladkrabang (KMITL), Bangkok, Thailand.

Strong crops inter. Co. Ltd. Samut Prakarn province, Thailand.

During of the Research

The research will be conducted from October 2014 to September 2015.

Chapter 2

Review of Literature

The living together of unlike organisms is known as symbiosis. . The term symbiosis (from the Greek: *syn* "with" and *biosis* "living") can play a roll in different kind of associations. In natural ecosystems plants are potential hosts for a broad spectrum of microbes (usually bacteria and fungi), the microbes that live on surface of plants call as epiphytes and microbes that live inside tissue of plant call as endophytes. They can be broadly categorized into mutualism, commensalism and parasitism according to benefits for both partners (mutualistic symbiosis), microbial benefit not affecting the host plant (commensalistic symbiosis) or negative impact on host fitness (pathogenic or parasitic interaction) (Paszkowski, 2006). Endophytes are organisms that live inside the plant without apparent any symptom of infection (Gimenez, *et al.* 2007).

2.1 Endophytic fungi

2.1.1 Definition and classification of endophytic fungi

“Endophyte” was named by Debarry (1866) to describe fungi ranging from virulent foliar pathogens to mycorrhizal root symbionts. The term endophyte comes from the Greek (*endo* “within” and *phyton* “plant”) that include a broad spectrum of plant endosymbionts from bacteria to insects colonizing inside any organ of the plant with variable life styles (Schulz and Boyle, 2005). Endophytes include an assemblage of microorganisms with different life style: some endophyte perform as saprophyte on dead or senescing tissue, mutualisms that can protect the host plant from pathogen or can promote growth the host plant, but also can be the latent pathogens and virulent pathogens at early stages of infection. These parasitic interactions may vary from mutualistic to commensalistic to latently pathogenic and exploitive. Phenotypes of the interactions are often plastic, depending on the genetic dispositions of the two partners, their developmental stage and nutritional status, but also on environmental factors (Schulz and Boyle, 2006).

Fungal endophytes are fungi that grow inside living plant tissues for at least part of their life cycle without causing disease symptoms (Stone *et al.*, 2004). There is some debate over where the boundaries of this definition should exist (Schulz and Boyle, 2005). Generally, an endophytic relationship refers to a mutualistic relationship with a positive impact on the fitness of both organisms; however, antagonisms in species-specific interactions have been demonstrated (Saikkonen *et al.*, 2004, 2006).

Based on phylogeny and life history traits, endophytic fungi were originally organized into two broad groups – clavicipitaceous endophytes (CEs) colonize shoots and rhizomes of a narrow host range of cool- and warm-season grasses (*Poaceae*) and non-clavicipitaceous endophytes (NCEs) have been isolated from shoots and/or roots of almost all sampled plants and are plentifully diverse. More recently, these groups have been divided into four distinct functional classes based on a number of different criterion related to life history. The classification into one of the four classes based on: host range (single species to highly ubiquitous), tissue colonization (localized to general), extent in plant colonization (limited to extensive) and biodiversity (low to high), mode of transmission (vertical to horizontal) and host, most of them belonging to Ascomycota. (Rodriguez *et al.*, 2009).

The CEs (class 1 endophytes) belong to the fungal teleomorphic genera *Epichloë* and *Balansia* correspond to the anamorphs *Neotyphodium* and *Ephelis*, respectively. They form systemic intercellular infections with a hyphal gradient along the plant axis and primarily transmitted vertically by seeds (Schulz and Boyle, 2005; Kuldau and Bacon, 2008).

The NCEs have been divided into three functional groups, class 2 class 3 and class 4. Class 2 endophytes broadly colonize both above and below ground tissues, but with limited biodiversity in individual plant hosts. They are both transmitted vertically and horizontally, sometime colonize non-habitat or habitat adapted benefits to the plant host. Endophytic fungi in this class belong to a few

This material is reserved for educational use only; not allowed for commercial use.

members of the *Agaricomycotina*, *Pucciniomycotina* *Basidiomycota* and also to *Pezizomycotina* (Ascomycota) such as *Phoma* sp., *Arthrotrichum* sp., *Fusarium culmorum*, *Colletrichum* sp. and *Curvularia protuberata* that can also found in roots, rhizomes, stems and leaves (Rodriguez *et al.*, 2009, Llorca *et al.*, 2006).

NCEs class 3 the colonization has restricted to shoots, but with high biodiversity in individual tissues. They have been isolated broadly from tropical forest to boreal and Arctic plant populations. They are horizontally transmitted by wind, rain and insects (Higgins *et al.*, 2007; Arnold, 2008; Feldman *et al.*, 2008). The Ascomycota that isolated from leaves the most belonging to the *Sordariomycetes*, *Dothidiomycetes*, *Pezizomycetes*, *Leotiomyces* and *Eurotiomyces*, while *basidiomycetous* isolates belonging to *Agaricomycotina*, *Pucciniomycotina*. Their benefits to the plant seem to be more complex and non-habitat-adapted (Arnold *et al.*, 2007; Higgins *et al.*, 2007; Rodriguez *et al.*, 2009., Vega *et al.*, 2010).

Class 4 endophytic fungi colonize only roots. They also known as Dark Septate Endophytes (DSEs), are easily distinguished by their highly melanized septate hyphae. These fungi are limited to the roots of their host, coexisting with mycorrhizal fungi but not growing outward into the rhizosphere. They have a broad host range and belong to different phylogenetic groups among Ascomycota and among non-mycorrhizal members of the order *Sebaciales* (Basidiomycota) (Weiss *et al.*, 2004; Addy *et al.*, 2005; Selosse *et al.*, 2009; Schäfer and Kogel, 2009).

2.1.2 Ecology of fungal endophyte

Fungal endophyte species are found across diverse habitats in the majority of plant species (Vega, 2008). They can reside in root, stem, leaf or multiple tissues (Stone *et al.*, 2006; Yuan *et al.*, 2009). As refer to many study of their diversity in many plants as follow.

Dandu Anitha *et al* (2013) isolated endophytic fungi from 5 medicinal plants viz *Boswellia ovalifoliolata*, *Pterocarpus santalinus*, *Shorea thumbuggaia*, *Syzygium alternifolium*. 14 fungal species, viz., *Fusarium oxysporum*, *Colletotrichum falcatum*, *Pestalotiopsis* sp. , *Aspergillus fumigatus*, *Aspergillus flavipes*, *Sterile mycelia*, *Penicillium senticosum*, *Gliocladium roseum*, *Phomopsis jacquiniana*, *Nigrospora sphaerica*, *Leptosphaeria* sp., *Phomopsis archeri* , *Alternaria alternata*, *Aspergillus niger* were isolated. They found that fungi were most diverse in leaves and root while in stems have low diverse of endophytic fungi. In opposite of this research and these plants Sun Y. *et el.* (2011) isolated endophytic fungi from *Suaeda* species, annual halophytes growing in soils with high salinity and high concentration of irons where non-halophytes are unable to thrive, found *Alternaria* sp., Ascomycete sp., *Bipolaris* sp., *Camarosporium* sp., *Cladosporium* sp., *Fusarium* sp., *Penicillium* sp. and *Phoma* sp. the isolation frequency in stems was higher than in leaves.

Glenn A. and Bodri MS. (2012) isolated endophytic fungi from 4 species of the carnivorous pitcher plant genus *Sarracenia*: *S. minor*, *S. oreophila*, *S. purpurea*, and *S. psittacina*. Twelve taxa of fungi, 8 within the Ascomycota and 4 within the Basidiomycota, were identified. They speculate that endophyte-infected *Sarracenia* may benefit from their fungal associates by their influence on nutrient availability from within pitchers and, possibly, by directly influencing the biota within pitchers.

Sun Y. *et al.* (2011) in China isolated endophytic fungi from the stems and leaves of *Suaeda* sp. found 2 species i.e. *Alternaria* sp. and *Phoma* sp.

Saithong P. *et al.* (2010) from Thailand isolated the endophytic fungi from the bark of *Cephalotaxus mannii* (plum-yew) found 13 genera, viz. *Cladosporium* sp., *Acremonium* sp., *Trichoderma* sp., *Monilia* sp., *Fusarium* sp., *Spicaria* sp., *Humicola* sp., *Rhizoctonia* sp., *Cephalosporium* sp., *Botrytis* sp., *Penicillium* sp., *Chalaropsis* sp. and *Geotrichum* sp.

Kumar, D.S.S. and Hyde, K.D. (2004) reported that 60 taxa including 30 morphotypes were isolated from the different parts of the Chinese medicinal plant, *Tripterygium wilfordii*. The endophytic assemblages of *T. wilfordii* comprised a number of cosmopolitan species such as *Colletotrichum gloeosporioides*, *Guignardia* sp., *Glomerella cingulata*, *Pestalotiopsis* sp., *Phomopsis* sp. and *Phyllosticta* sp. The overall fungal community of *T. wilfordii* was moderately diverse. The fungal community from the twig xylem parts was most diverse, followed by leaves, twig bark, root xylem and flowers.

As follow study we can indicate that endophytic fungi ecology are very diverse, but there is a question that how can endophytic fungi colonize in plant tissue from one generation to one generation. About this question Yuan *et al.* (2009) conclude that endophytic fungi transfer from each plants by 2 ways. The first one is vertical transfer that direct transfer by seeds. The second is horizontal transfer that can transfer by air, rain and insect in environment who have high taxonomic and host diversity, and can broadly colonize all plants in an eco-system.

2.1.3 Secondary metabolites produced by edophytic fungi

Endophytes may produce new bioactive substances because of a continual metabolic interaction with their hosts and the environment and this metabolic interaction is important for the symbiosis of both partners (Schulz and Boyle, 2005; Suryanarayanan *et al.*, 2009). Many researchers try to indicate for antimicrobial from many plants. Below there are many research about isolate antimicrobial substance from endophytic fungi.

From China Su Z-Z *et al.* (2013) investigated the dark septate endophyte (DSE) *Harpophora oryzae* in rice roots and its biocontrol potential in rice blast disease caused by *Magnaporthe oryzae*. The result showed that the colonization pattern of *H. oryzae* was consistent with the typical characteristics of DSEs. *H. oryzae* enhanced local resistance by reactive oxygen species (ROS) and high antioxidative level and induced OsWRKY45-dependent SA-mediated systemic resistance against rice blast. Yin Lu *et al.* (2012) isolated Endophytic fungi from *actinidia macrosporma* investigated for their bioactivity indicated that fungi from this plant exhibited toxicity against brine shrimp and antitumour. Liang H. *et al.* (2012) indicated that endophytic fungi from ornamental plant like *Ophiopogon japonicus* (Liliaceae) reported that this plants also produce metabolite inhibited *Staphylococcus aureus* and *Cryptococcus neoformans*. And in previous year Gao Y. *et al.* (2011) reported that 4 strains of endophytic fungi from roots of Pigeon Pea (*Cajanus cajan* (L.) Millsp.) were producing Cajaninstilbene acid (CSA, 3-hydroxy-4-prenyl-5-methoxystilbene-2-carboxylic acid), CSA is responsible for the prominent pharmacological activities in pigeon pea. The five endophytic isolates were characterized by analyzing the internal transcribed spacer (ITS) rRNA and b-tubulin genes. The 4 strains isolated from pigeon pea roots were found to be closely related to the species *Fusarium oxysporum* and another one was identified as *Neonectria macrodidym*.

While in Philippine Jeremy Martin O. Torres and Thomas Edison E. dela Cruz (n.d.) indicate that crude isolated endophytic fungi from *Canarium ovatum* show that *Colletotrichum* sp. PFE31 crude extract exhibited broad-spectrum antibacterial activity against representative Gram-positive, Gram-negative, and spore-forming bacilli. Melfei E. Bungihan *et al.* (2013) isolated Endophytic fungi from *Pandanus amaryllifolius*. The genera *Colletotrichum*, *Chaetomium*, *Diaporthe*, *Glomerella*, *Guignardia*, *Lasiodiplodia*, *Lulworthia*, *Phoma*, *Phyllosticta*, *Trichoderma*, and *Truncatella*. were identified by morphology. The selected species was use for antimicrobial, show that can against *Mycobacterium tuberculosis*, *Staphylococcus aureus*, *Eschericia coli*, and *Gordonia terrae*.

Oil-Seed Crop *Jatropha curcas* was selected for endophytic fungi in India by Susheel Kumar and Nutan Kaushik (2013). *Colletotrichum truncatum*, *Nigrospora oryzae*, *Fusarium proliferatum*, *Guignardia cammillae*, *Alternaria destruens*, and *Chaetomium* sp. were selected from leaf of

This material is reserved for educational use only, not allowed for commercial use.

Jatropha curcas. Dual plate culture bioassays and bioactivity assays of solvent extracts of fungal mycelia showed that isolates of *Colletotrichum truncatum* were effective against plant pathogenic fungi *Fusarium oxysporum* and *Sclerotinia sclerotiorum*. Susheel Kumar, Nutan Kaushik and Peter Proksch (2013) isolated *Chaetomium globosum* from *Withania somnifera*. They used Ethyl acetate, hexane and methanol extracts method to show the control of *Chaetomium globosum* against *Sclerotinia sclerotiorum*. The result showed that Ethyl acetate and methanol extracts were more effective than hexane extract. Qadri M. *et al.* (2013) reported that endophytic fungi of selected plants from the Western Himalayas, India. Seventy two strains of endophytic fungi were isolated. Several strains of these endophytic fungi were showed that against *E. coli* and *S. aureus*. 24 endophytes inhibited three or more phytopathogens 17 fungi possessed immuno-modulatory activities with five of them showing significant immune suppression as demonstrated by the in vitro lymphocyte proliferation assay. Kumar A. *et al.* (2013) isolated endophytic fungi from *Catharanthus roseus* plant and found a fungus that can produce vinblastine and vincristine, anti-cancer drugs, in appreciable amounts. That endophytic fungi have identified as *Fusarium oxysporum*.

In Thailand Tiamphet M. and Nuangmek W. (2013) isolated endophytic fungi from Eucalyptus and another 29 species of herbs. Two hundred and seventy endophytic fungi were isolated. Unknown, *Phomopsis* sp. and *Xylaria* sp. were the most common species. Dual culture method was used to screening for evaluation on the potential effectiveness of candidate biological control agents against rot disease of eucalyptus (K7) seedling caused by *Rhizoctonia solani*. Revealed that *Nodulisporium* sp. and *Muscodor* sp. were strong anti-fungal activity against *R. solani* and also have good result in green house. In the same year in southern Thailand Supaphon P *et al.* (2013) reported that seven isolates endophytic fungi from three seagrasses species, *Cymodocea serrulata*, *Halophila ovalis* and *Thalassia hemprichii* including *Hypocreales* sp. PSU-ES26 from *C. serrulata*, *Trichoderma* sp. PSU-ES8 and PSU-ES38 from *H. ovalis*, and *Penicillium* sp. PSU-ES43, *Fusarium* sp. PSU-ES73, *Stephanonectria* sp. PSU-ES172 and an unidentified endophyte PSU-ES190 from *T. hemprichii* exhibited strong antimicrobial activity against human pathogens with minimum inhibitory concentrations (MIC) of less than 10 µg/ml. The inhibitory extracts at concentrations of 4 times their MIC destroyed the targeted cells as observed by scanning electron microscopy. These results showed the antimicrobial potential of extracts from endophytic fungi from seagrasses.

In Taiwan Min Y.H. *et al.* (2012) isolated endophytic fungi from medical herbs of *Lauraceae* and *Rutaceae*. 156 isolates of endophytic fungi collected from twigs of medicinal plants of *Lauraceae* (67 isolates) and *Rutaceae* (89 isolates). The most common endophytes were in the taxa of *Colletotrichum*, *Guignardia*, *Hypoxylon*, *Nigrospora*, *Phomopsis* and *Xylaria*, and the most common hosts were *Citrus* and *Zanthoxylum* of *Rutaceae* and *Cinnamomum* of *Lauraceae*. Endophytic fungi of *Lasmenia* sp. Isolate significantly reduced severity of anthracnose of Chinese cabbage caused by *Colletotrichum higginsianum* under greenhouse conditions.

In Korea Paul N. C. *et al.* (2011) reported that a total of 481 isolates were recovered from the surface sterilized tissues of leaves, stems and roots of chili pepper (*Capsicum annuum* L.) plants in seedling, flowering and fruiting stages. Twenty one fungal genera were characterized, belonging to 16 Ascomycota and 5 Basidiomycota. *Penicillium* in seedling stage, *Fusarium* in flowering stage, *Colletotrichum* followed by *Fusarium*, *Alternaria* and *Xylaria* in fruiting stage was predominant and *Alternaria*, *Cladosporium* and *Fusarium* were common in all growth stages. Ninety phenotypes were evaluated for the antimicrobial activity against three major pathogens (*Phytophthora capsici*, *Colletotrichum acutatum* and *Fusarium oxysporum*) of chili pepper. Among them 16 isolates inhibited the growth of at least one test microorganisms. Three strains showed a broad spectrum antifungal activity and displayed strong inhibition against chili pepper pathogenic fungi.

In Brazil Hanada R.E. *et al.* (2010) isolated endophytic fungi from stems and branches of *Theobroma cacao* (cacao) and *Theobroma grandiflorum* trees growing in the Amazon region of Brazil. One hundred and three isolates that could be classified at least up to genus level were tested

This material is reserved for educational use only, not allowed for commercial use.

Forbidden to modify the content, and cite the document when use. ศราวุฒ พลโพธิ์ วรณะ สัว

against *P. palmivora* in pods attached to cacao trees in the field. Results indicated that 70 % of isolates showed biocontrol effects to a certain extent, suggesting that culturable endophytic fungal biodiversity in this system is of a mostly mutualistic type of interaction with the host.

In Panama of USA Herre A.D. *et al.* (2007) isolated endophytic fungi from *Theobroma cacao* and showed that the endophytic fungi against pathogen (*Phytophthora palmivora*).

Endophytic fungi also can promote plant growth to growing up better. They are able to produce phytohormones such as Gibberelin (GAs) and indoleacetic acid (IAA) and mitigate abiotic stresses like salinity and drought. Some research of plant growth promotion of endophytic fungus as follow.

Rola S. Mahmoud, Kazuhiko Narisawa (2013) Isolated endophytic fungi from roots of tomatoes and cabbages. Two hundred and five isolates have showed that the most fungi dominant by *Fusarium sp.* and a new fungal endophyte, *Scolecobasidium humicola*, was identified as a common dark septate endophytic fungal (DSE) species under both natural and agricultural conditions. This fungus was found to grow endophytically in the roots of tomato seedlings. Two isolates of *S. humicola* have shown the ability to increase plant biomass with an organic nitrogen source. This finding is the first report of *S. humicola* as an endophyte and could help to improve plant growth with organic nitrogen sources.

Waqas M. *et al.* (2012) isolated endophytic fungi from the root of cucumber. 2 strains of fungal endophyte, *Phoma glomerata* LWL2 and *Penicillium sp.* LWL3, were selected and further to examine for their production of gibberelins (GAs) and indoleacetic acid (IAA). The result showed that both fungi strain can promote shoot length of two rice varieties, *Waito-C* and *Dongjin-beyo* rice. The cultures of *P. glomerata* and *Penicillium sp.* also contained IAA. The culture application and endophytic-association with host-cucumber plants increased the plant biomass and related growth parameters under sodium chloride and polyethylene glycol induced salinity and drought stress as compared to control plants. The endophytic symbiosis resulted in higher assimilation of essential nutrients like potassium, calcium and magnesium as compared to control plants during salinity stress. Endophytic-association reduced the sodium toxicity and promoted the host-benefit ratio in cucumber plants as compared to non-inoculated control plants. The symbiotic-association mitigated stress by compromising the activities of reduced glutathione, catalase, peroxidase and polyphenol oxidase. Under stress conditions, the endophyte-infection significantly modulated stress through down-regulated abscisic acid, altered jasmonic acid, and elevated salicylic acid contents as compared to control. In conclusion, the two endophytes significantly reprogrammed the growth of host plants during stress conditions.

Khan S.A. *et al.* (2012) investigated plant growth promoting activity of roots inhabiting endophytic fungi in order to evaluate their role in the survival of host plants under extreme sand dune environment of coastal regions. One hundred and twenty two fungal isolates were collected from the roots of 9 sand dune plants and were screened for growth promoting secondary metabolites. The results showed that 101 fungal isolates (82.7%) promoted plant height and shoot length of *Waito-C* rice, while 21 fungal isolates (17.2%) inhibited growth attributes. The fungal isolate *Gibberella fujikuroi* along with distilled water and Czapek broth medium were used as control during the experiment. It was concluded that a major proportion of endophytic fungi inhabiting sand dune plants produce metabolites, which are helpful in plant growth and development.

Hamayun M. *et al.* (2010) isolated endophytic fungi from cucumber (*Cucumis sativus* L.) found 19 strains of fungi. Pure cultures of 19 endophytic fungi were tested for shoot length promotion of *Waito-C* rice to identify the GA production capacity of these fungal isolates. One of these isolate showed that increased shoot length of *Waito-C*, in comparison to control treatments.

This material is reserved for educational use only, not allowed for commercial use.

Forbidden to modify the content, and cite the document when use. ศราวุฒ พลโพธิ์ วรณะ ส้ว

2.1.4 Host benefit of infection endophytic fungi

Fungal endophytes, and their effects on plant metabolism, play a large role in phenotypic plasticity and environmental adaptability and likely played a unique role in selection and speciation in many host species (Rudgers *et al.*, 2009). Plant can supply nutrients to endophyte, and endophyte will release active metabolites through metabolic activities. Those active metabolites can enhance the roots development and assist solubilizing phosphorus, with which would increase the ability of plants to absorb nutrients from the soil (Li, J.H., *et al.*, 2008). Because of those activities, endophyte can promote host growth and supply resistance mechanism of host. As the study of Dai C.C. *et al.* (2008) who inoculated the endophytic fungi *fusarium sp.* strain E4 and E5 that isolated from medicinal plant, *Euphorbia pekinensis*, to the host plant in pot culture. The result exhibited significant increase in growth and all tested growth parameters, as compared to the control.

There is a wide spread phenomenon about the symbiotic relationships between endophytes and plants. But this relationships can switch between mutualistic and parasitic (Kogel, K.-Het *al.* 2006). While a single gene changes can alter the mechanism of endophyte. For example, *E. festucae* which is a wild-type grows systemically in intercellular spaces, can promote the grown and enhance the resistance of plant, but while a single-copy plasmid of a NADPH oxidase gene, *noxA*, was inserted in the coding region, the plant would lose its apical dominance, become severely stunted, show precocious senescence, and eventually die (Tanaka, A., *et al.*, 2006). The same endophyte colonized in different host can react in the different lifestyles. *Colletotrichum magna* can cause anthracnose in cucurbit plants, but can grow asymptotically when colonized in various non-cucurbit species (Kogel, K.-Het *al.* 2006).

2.2 Orchids

Orchids are the monocotyledon taxonomy in Order Orchidales, Family Orchidaceae. There are 15000 to 25000 of orchid species distributed all around the world, mostly can found in tropical environment. They all can divide in 2 classifications as epiphytic orchids and terrestrial orchids. In Thailand found 500 genus and 800 orchid species. The special hereditary characteristic of orchid species is having many tiny seed in the pod and there are not growing and no endosperm in embryo. So to be germinated the orchids have to depend on some fungi species to be the energy and food sources such as many nutrients that important for growth (Thamasiri K., 2005).

Orchids are among the plant groups that have aroused most widespread interest among scientists and horticulturists, both for their study and use. They have been subject to particularly high commercial demand over the past 40 years for the beauty of their structure and their vividly colored flowers, steeped in symbolism and mystery. Some countries have declared orchid species as their national flowers (Clemente M., 2009). In Thailand orchids are the most valuable product for export and make money for country. Last years (2013) the export of orchids value 563,134,221 Bart, and the country that most import Thai orchids is China, Japan, U.S.A, India and Vietnam, respectively (Source: Office of Agriculture Economics). It takes many interested in planting orchid for commercial from businessman and horticulturists. So when there are plentiful culture of orchids that's mean it's following by the distributed of disease in 2012 . There are many study that compiled the pathogen fungus on orchids include *Alternaria sp.*, *Botrytis cineria*, *Cercospora sp.*, *Cladosporium sp.*, *Colletotrichum gloeosporioides*, *Curvularia eragrostidis*, *Diplodiasp.*, *Drechslera sp.*, *Fusarium sp.*, *Nectria sp.*, *Phyllosticta pyriformis*, *Pseudocercospora dendrobii*, *Phytophthora nicotiana* var. *parasitica*, *Phytophthora palmivora*, *Rhizotonia sp.*, *Schlerotium rolfsii* and *Volutela sp.* (Thamasiri K., 2005; Sutthiwari P., 2006).

2.3 Previous study about endophytic fungi in orchids

Juan Chen *et al.* (2013) isolated endophytic fungi from 7 *dendrobium* sp. (*Orchidaceae*) to search for *Xylariaceae* fungi. The result showed that A total of 961 culturable endophytic fungi were isolated from seven species of *Dendrobium*. In the original 961 cultures, the xylariaceae taxa had the highest isolation frequency, 22.58% (217/961), followed by *Fusarium* at 10.71% (103/961), *Colletotrichum* at 6.55% (63/961), and *Phomopsis* at 4.37% (42/961).

Sawmya K. *et al.* (2013) compared the leaf and root endophyte assemblages of two orchids (*Bulbophyllum neilgherrense* and *Pholidota pallida*) from natural forests and greenhouse conditions. The result found that *Xylariaceae* sp. were consistently associated with leaf and root tissues, while *Guignardia* and *Pestalotiopsis* were found predominantly in the leaf tissues of both orchids. The endophyte assemblages showed that the endophytes exhibited distinct organ but little host specificity. Endophytic fungi have more divers in green house condition more than in forest condition. And the roots showed more diverse than in leaves.

Mangunwardoyo W, Suciati and Gandjar I. (2012) were isolated endophytic fungi from *Dendrobium crumenatum* (Pigeon orchid) and investigated antimicrobial activity against 5 pathogens namely *Candida albicans* ATCC 2091, *Candida tropicalis* LIPIMC 203, *Escherichia coli* ATCC 25922, *Bacillus subtilis* ATCC 6633 and *Staphylococcus aureus* ATCC 25923. The result showed that twelve species of endophytic fungi were identified from 60 samples obtained from *D. crumenatum*. namely *Cladosporium cladosporioides*, *C. sphaerospermum*, *Colletotrichum gloeosporioides*, *Colletotrichum* sp., *Curvularia brachyspora*, *Fusarium nivale*, *F. solani*, *Guignardia endophyllicola* (anamorf: *Phyllosticta capitalensis*), *Pestalotiopsis* sp., *Scolecobasidium* sp., *Westerdikella* sp., and *Xylohypha* sp. The dominant endophytic fungi was *G. endophyllicola*. For investigated the antimicrobial showed that *Fusarium nivale* was able to inhibit the growth of *C. albicans* and *C. tropicalis*. However, others specimens did not inhibit the tested bacterial, which are Gram-negative *E. coli* and Gram-positive *B. subtilis* and *S. aureus*.

Kasmir J. *et al.* (2011) isolated endophytic fungi from 4 species epiphytic orchids namely *Bulbophyllum kaitiense* Reichebt. *Gastrochilus acaulis* (Lindley) Kuntze *Dendrobium nanum* Hook.f and *Geodorum densiflorum* (Lam). Schltr. The two important fungi belonging to tricomaceae members *Aspergillus terreus* and *Penicillium aculeatum*.

Tempesta S. *et al.* (2011) isolated endophytic fungi to screening for *Pestalotiopsis* endophyte from leaves of two orchid species: Among the fungi isolates from orchids 29 strains of *Pestalotiopsis* sp. was discovered such as *P. sydowiana*, *P. photiniae*, *P. leucothoes*, *P. clavispora*, *P. crassiuscula*, *P. rhododendri*, *P. versicolor*, *P. photiniae*, *P. gracilis*, *P. aquatic*, *P. palmarum*, *P. theae*, *P. maculans*, *P. funerea*, *P. neglecta*, *P. disseminata*, *P. adusta*, *P. microspore*, *P. bacilia* and *P. vismiae*.

Valdes M. *et al.* (2011) investigated fungi from root of terrestrial orchid *Cypripedium irapeanum* found 10 isolates 8 strains of fungi : 3 identified as endophytic fungi, *Fusarium* sp. and *Cylindrocarpon* sp., 3 as micorhiza, *Sistotrema* sp., *Rhizoctonia solani* and *Epulorhiza* sp., 3 as dark septate endophytes, MUT 885 and *Phomopsis* sp., 1 as biocontrol fungus, *Gliocladium catenulatum*.

Juntawong J. *et al.* (2010) investigated endophytic fungi from terrestrial orchids *Geodorum* roots in three different seasons in deciduous forest in northern Thailand. The result showed that 198 isolates comprising to 13 genera. The major genera were *Tricoderma* sp. followed by *Fusarium* sp., the least were *Aspergillus* sp., *Colletotrichum* sp., *Eupenicillium* sp. and *Helicoma* sp. In rainy season showed high diversity of fungi more than dry and cold season. This study indicated that changes of physical factors in different seasons had effect on endophytic fungi diversity.

Gezgin Y. and R. Eltem (2009) were investigated in orchids from the Aegean and Mediterranean regions. Four species of orchids as follow *Anacamptis pyramidalis* (L.) L.C.M.Richard, *Orchis sancta* L., *Ophrys fusca* Link., and *Serapias vomeracea* subsp. *orientalis*

This material is reserved for educational use only, not allowed for commercial use.

Forbidden to modify the content, and cite the document when use. ศราวุฒ พลโพธิ์ วรณะ ส้ว

Greuter were isolated for endophytic fungi. The result showed that a total of 47 isolates, having genus characterisations as 44 (94%) isolates belonging to the genus *Fusarium*, 2 (4%) isolates belonging to the *Rhizoctonia* DC. ex Fr.-like fungi, and 1 (2%) isolate belonging to the genus *Papulaspora* Preuss, were found from the orchid root and tubers.

Tao G. *et al.* (2008) investigated endophytic fungi from terrestrial orchid *Bletilla ochracea*. There were many species of fungi have revealed in this study. Fourteen species were isolated from the leaves, *Alternaria* sp., *Ascomycete* sp., *Chaetothyriales* sp., *Cladosporium* sp., *Colletotrichum* sp., *Cryptococcus* sp., *Dioszegia* sp., *Epulorhiza* sp., *Gibberella* sp., *Herpotrichiellaceae* sp., *Leptosphaerulina* sp., *Mycosphaerella* sp., *Pestalotiopsis* sp. and *Phaeosphaeria* sp. Nine species were isolated from roots, *Cercospora* sp., *Cylindrocarpon* sp., *Exophiala* sp., *Nectria* sp., *Fusarium* sp., *Gibberella* sp., *Herpotrichiellaceae* sp., *Neonectria* sp. and *Sebacina* sp. As the result of this study micorrhiza fungi were isolated from leaves. The diversity within leaves was higher than that within roots. Fungal communities within leaf and root tissues were significantly different.

Ovando I. *et al.* (2005) screened endophytic fungi in roots of mature plants of epiphytic orchid *Cattleya skinneri* to test their potential to improve seed germination in *C. skinneri* and develop in acclimatizing plantlets of *C. aurantiaca* and *Brassavola nodosa*. One hundred and eight isolates belonging to 11 genera consist of *Tricodermasp.*, *Verticilliumsp.*, *Aspergillussp.*, *Epulorhizasp.*, *Fusarium* sp., *Monilliopsis* sp., *Pestalotiopsis* sp., *Botrytis* sp., *Penicilliumsp.*, *Penicillifera* sp. and *Tetracladium* sp. Eight strains in this result used for promote seed germination of *C. skinneri* and 3 strains used for promote plant growth of *B. nodosa*. None of these strains can promote both of experiments.

Bougoure J.J. and J.D.W. Dearnaly (2005) isolated endophytic fungi from roots of *Dipodium variegatum* found *Russula occidentalis*, *R. salaris*, *R. lepida*, *Verticilliumsp.*, *Tricoderma hamatum*, *Halocyphina villosa* and *Merismodes fasciculata*.

Chapter 3

Materials and Methods

The sample will be collected in greenhouse condition from Rajamangala University Technology Tawan-ok Chantaburi Campus, Chantaburi province, Thailand. Samples of Leaves and roots from 9 varieties of selected orchids will be packed in sterile polythene bags to keep moisture and bring to laboratory over 12 hours after collecting. The name of nine orchid varieties will be collected as follows:

1. *Grammatophyllum scriptum*
2. *Cymbidium dayanum* Rchb.F
3. *Dendrobium hercoglossum* Rchb.f.
4. *Dendrobium palpebrae* Lindl.
5. *Torenia fournieri* Lindl. ex Fourn.
6. *Doritis pulcherrima* var. *buyssonianana*
7. *Dendrobium crumenatum* Sw.
8. *Dendrobium friedericksianum* Rchb.f.
9. *Grammatophyllum specinocum* BL.

2. Isolation of endophytic fungi

The surface sterilization will be done following the procedure used by Helander *et al.* (2007) and Blumenstein (2010) with some modifications

- Cut approximately 0.5 cm² long piece from each leaf and root of orchids; will try to remove epidermal tissues gently by razor blade from leaves and remove outer cuticle from root sections too.
- Thoroughly washed in sterile water to remove dirt;
- Dip the samples into 75% ethanol for 3 minutes;
- Dip the samples in 0.5% Na-hypochlorite solution for 2 minutes;
- Wash the samples in sterile water for 3 times;
- Dry the samples in sterile tissue paper until dehydrate.

The procedure will be carried out under a laminar hood in aseptic condition. After removing the excess water, the segments were placed on Petri plates (9 cm diam.) containing with WA (water agar). The plates were incubated at room temperature and fungal colonies will be regularly observed. The Petri dishes will be then periodically checked every day for fungal growth. Each time a seen new colony from inner leaf tissue and root segments, it will be sub-cultured on PDA (Potato Dextrose Agar) plates to make pure cultures.

3. Identification of endophytic fungi by morphological and physiological characterization


For characterization of the morphology of fungal isolates, slides will be prepared from cultures and will be stained with bromothymol blue reagent in lacto phenol and will be examined with a binocular compound microscope. Identification will be based on morphological characteristics such as growth pattern, colony and hyphae, colour of colony and medium, surface texture, margin character, aerial mycelium, mechanism of spore production and characteristics and size of the spore (H. L. Barnett and B. B. Hunter, 1956).

The morphological characteristics will also be followed Crous *et al.* (2009) that used as criterion to the group all fungi. Each fungus will be observed and examined to the characteristics and will be assigned to a specific morphological group. To be part of the same group, a fungus will be grouped.

This material is reserved for educational use only, not allowed for commercial use.

Forbidden to modify the content, and cite the document when use. ศรายุทธ ผลโพธิ์ วรรณะ ต้ว

- Colour. The colony colour will be observed to discriminate the fungi. If a fungal colony will be seen more than one distinct colour, the description will be done from the middle towards the outer part.
- Agar colour. Some isolates may possibly be released metabolites or substances to change the colour of the agar where the colony is growing.
- Liquid drops. The presence and the colour of the liquid drops that some fungi produced on their surface will be recorded.
- Colony shapes. Each colony will be observed to grow in a certain way forming different shapes. The colony shapes will be distinct: crenate, entire edge, erose or dentate, fimbriate, lobate, radially striate with lobate edge, undulate, with rhizoids.
- Colony texture will be distinct as follow:- woolly, velvety, hilly, spongy, dry, furry and creamy.
- Spore characteristics will be noted. If the fungi would be developed a clear accumulation of spores on their surfaces and the colour of the spores will be observed.
- The growth rate will be observed as an additional characteristic in order to increase the common characteristic within a group.



Chapter 4

Results and Discussions

Many isolates can't identify the species because it doesn't produce spores. Twelve species belonging to Ascomycota were isolated from 9 orchid varieties. They are as follow:

1. *Chaetomium cochliodes*

Habitat: *Grammatophyllum scriptum*, *Cymbidium dayanum*, *Dendrobium hercoglossum* and *Torenia fournieri*.

The colonies are quite slowly growing, for cultured on PDA medium in room temperature for 7 days have 7 cm in diameter, cottony and white in color initially and then become soft-green. *C. cochliodes* has distinctive small brown 'lemon' or 'football' shaped ascospores. (Arx, J.A. von *et al.* 1986).

This species also isolated as endophytic fungi from *Cirsium arvense*, (Gange AC. *Et al.*, 2012). Li Gy *et al.* 2006 indicated that epipolythiodio-xopiperazines and chaetocochin were isolated from the ethyl acetate extract of the solid-state fermented rice culture of the fungus *Chaetomium cochliodes*. Their structures were elucidated on the basis of spectroscopic analysis. Compounds 1, 3, and 4 exhibited significant cytotoxicity *in vitro* against cancer cell lines Bre-04, Lu-04, and N-04,




Fig. 1: *Chaetomium cochliodes* A: The colony that culture on PDA 30 days, B The spores under microscopes 10X C The ascospores under microscopes 40X

2. *Chaetomium cupreum*

Habitat: *Grammatophyllum scriptum*.

The growth was quite slow, cultured on PDA medium for 7 days the growth have 4.4 cm in diameter. The hyphae are red and it produces the red color around mycelia.

Septate hyphae, perithecia, asci and ascospores are visualized. Perithecia are large, red color, fragile, globose to flask shaped and have filamentous, hair-like, brown to black appendages (setae) on their surface. Perithecia have ostioles (small rounded openings) and contain asci and ascospores inside (De Hoog *et al.*, 2000).

This species was isolated as endophytic fungus from *Macleaya cordata* and have ability to antifungal activity against *Rhizoctonia solani*, *Botrytis cinerea*, *Pytium multimum*, and antitumor activities against HL-60 and A549. It is toxic to *Artemiasalina* (Bi Z.M. *et al.* 2010).



Fig. 2: *Chaetomium cupreum* A: The colony cultured on PDA 30 days B: The spores under

This material is reserved for educational use only, not allowed for commercial use.

Forbidden to modify the content, and cite the document when use. ศราวุฒ พลโพธิ์ วรรณะ สัว

microscops 10X and C:Theascospores under microscops 40X

3. *Xylaria* sp.

Habitat: *Cymbidium dayanum*, *Dendrobiumhercoglossum*, *Doritispulcherrima*, *Dendrobiumcrumenatum*, *Dendrobiumfriedericksianum* and *Grammatophyllumspecinocum*.

The fungi have white mycelia; the growth was quite slow, for cultured on PDA in room temperature for 7 days have 5.8 cm in diameter. They produce the stroma on 20th day but didn't produce ascus and ascospore.

Xylaria sp. was the common species that can be isolated from other orchid varieties. Previous research also found that this species is most colonized in 7 *Dendrobium* sp. in China (Chen J. et al. 2013). Sawmya K. et al. (2013) also isolated *Xylaria* sp. from *Bulbophyllumneilgherrense* and *Pholidotapallida*.




Fig. 3: *Xylaria* sp. A: Colony culture on PDA 30 days B: Mycelia of *xylaria* sp. under microscops 40X

4. *Colletotrichum* sp.

Habitats: *Cymbidium dayanum*, *Dendrobiumhercoglossum*, *Toreniafournieri*, *Dendrobium crumenatum* and *Grammatophyllumspecinocum*.

The growth on PDA medium 7 days have 8.5-9 cm, the mycelia initially white-grey and then become black-brown. The conidia have no color with rods shape (Busarakham K., 2001).

Previous study from India showed that the endophytic fungus, *Colletotrichum gloeosporioides* was isolated from *Justiciagendarussa*, a medicinal plant has the ability to produce Taxol that is a potent anticancer drug used widely in the treatment of a variety of cancers (Gangadevi V. and Muthumary J., 2008).




Fig. 4: *Colletotrichum* sp. A: The colony culture on PDA 30 days and B: The spores under microscope 40X

5. *Nigrospora* sp.

Habitat: *Dendrobiumhercoglossum*

Nigrospora grows quite fast and produces woolly colonies on potato dextrose agar (PDA) at room temperature, on 7th days has 9 cm in diameter. Color of the colony is white initially and then becomes gray with black areas and turns to black eventually from both front and reverse. Spores (conidia) are produced singly on swollen urn-shaped conidiophores and are egg-shaped to flattened-spherical, black, and often have an equatorial colorless line or germ slit.

This species also can isolated from deep sea invironment at Arabian Sea and The secondary metabolites produced from this organism showed potent antimicrobial and anticancer activities with immediate application to cosmetics and pharmaceutical industries. (Arumugam GK, *et al*, 2015)




Fig. 5: *Nigrospora* sp. A: The colony culture on PDA 30 days B: The hyphae and spores under microscope 40X

6. *Phoma* sp.

Habitats: *Cymbidium dayanum*, *Doritis pulcherrima* and *Grammatophyllum specinocum*.

The hyphae were growing rapidly, for 2 days that cultured on PDA medium have 6.5 cm in diameter. The hyphae first growth have white color and than become black-brown. Spores are colorless and unicellular. The pycnidia are black and depressed in the tissues of the host.

Phoma sp. are endophytic fungi that can isolated from many plants. Wang LW *et al.* (2012) isolated *phoma* sp. from chinese medicinal plant *Arisaema rubescens* and showed that the bioactive metabolites from this species can inhibited the growth of four plant pathogenic fungi (*Fusarium oxysporium*, *Rhizoctonia solani*, *Colletotrichum gloeosporioides*, and *Magnaporthe oryzae*). This species also isolated from roots of cucumber in Korea and showed that have ability to produce Gibberellins and Indoleacetic Acid and Promotes Host-Plant Growth during Stress (Waqas M. *et al.*, 2012).




Fig. 6: *Phoma* sp1. A: The colony culture on PDA 30 days and B: The spores and mycelia under microscope 40X

7. *Curvularia* sp.

Habitats: *Grammatophyllum scriptum* and *Doritis pulcherrima*.

The growth was quit fast for cultured on PDA medium for 2 days have 5.2cm in diameter. From the front, the color of the colony is white to pinkish gray initially and turns to black as the colony matures. From the reverse, it is dark brown to black. Conidia are pale brown, with three or more transverse septa (phragmoconidia) and are formed apically through a pore (poroconidia) in a sympodially elongating geniculate conidiophore similar to *Drechslera*. Conidia are cylindrical or slightly curved, with one of the central cells being larger and darker. Germination is bipolar and some species may have a prominent hilum (McGinnis, M.R. 1980).

Zakaria L. *et al.* (2010) isolated this species from *Oryza sativa*. Aharwal R.P. *et al.* (2014) also isolated this fungus from *Calotropis procera* (Linn.) R.Br. a widely used medicinal plant in India.




Fig. 7: *Curvularia* sp. **A:** The colony culture on PDA 30 days and **B:** The spores under microscope 40X

9. *Pestalotiopsis* sp.

Habitat: *Dendrobiumfriedericksianum* , *Cymbidium dayanum* and *Grammatophyllum scriptum*.

The mycelia are white in color. This species have quit slow of growing. For growing on PDA in room temperature for 7 days have 5.5cm in diameter. Conidiophores (annellides) produced within compact fruiting structures (aecervuli or pycnidia). Spores (conidia) 4 to 5 celled, with the two or three central cells dark brown, and with two or more apical appendages or hairs.

Pestalotiopsis sp. is the common endophytic fungi that can isolate from orchids and other plants. They also isolated from other orchid varieties from other country such as Tempesta S. *et al.* (2011) isolated endophytic fungi from leaves of *C. mrcranthum* and *O. adendrobium* also found 29 strains of *Pestalotiopsis* sp. Wei J. *Get al.* (2007) isolated this species that associated with plants of Podocarpaceae, Theaceae and Taxaceae in southern China also obtained 24 species. Kumaran R *Set al.* (2010) isolated *Pestalotiopsisver-sicolor* and *Pestaloti-opsisneglecta* as endophytic fungi from Japanese Yew tree, *Taxuscuspidata* and indicated that they can produce taxol concentration induces increased cell death in cancer.




Fig. 9: *Pestalotiopsis* sp. **A:** The colony culture on PDA 30 days **B:** The conidia under microscops 40X

10. *Corynascus* sp.

Habitat: *Cymbidium dayanum*

The fungus have quit fast in growth. Only 3 days the mycelia have 9.0 cm in diameter. It has black-white mycelia, produce lots of black spores and inside spores there are lots of little ascospores.




Fig. 10: *Corynascus* sp. **A:** The colony culture on PDA 30 days and **B:** The spores and acoospores under microscops 40X

11. *Cladosporium*-like fungi

Habitat: *Grammatophyllum scriptum*, *Cymbidium dayanum*, *Dendrobium palpebrae*, *Doritis pulcherrima* and *Dendrobium crumenatum*

The initial growth the mycelia have white color and then become black. The growth was quite fast. On 6th day of culture on PDA have 9 cm in diameter. The mycelia have branches and the lots of small spore.




Fig. 11: *Cladosporium*-like fungi A: The colony culture on PDA 30 days and B: The spores and mycelia under microscopes 40X

12. *Achaetomium* sp.

Habitat: *Grammatophyllum scriptum*

The mycelia growth look like the yellow sheets and this fungus growth quite slow, cultured on PDA medium in room temperature for 7 days have 5.5 cm in diameter. The spores are large with yellow in color. Inside the spores have many ascospores.



Fig. 12: *Achaetomium* sp. A: The colony culture on PDA 30 days B: The spores under microscopes 10X and C: The ascospores under microscopes 40X

Reference

- Agrios, G.N. (2005). Plant Pathology. Elsevier Academic Press. pp.922.
- Arnold, A.E., Henk, D.A., Eells, R.L., Lutzoni, F., and Vilgalys, R. (2007). Diversity and phylogenetic affinities of foliar fungal endophytes in loblolly pine inferred by culturing and environmental PCR. *Mycologia* 99:185–206.
- Arnold, A.E. (2008). Endophytic fungi: hidden components of tropical community ecology. In: Schnitzer S, Carson W, eds. *Tropical forest community ecology*. Malden, MA, USA: Blackwell Scientific, Inc. pp: 254-271.
- Anu Rajan, S.(2012). Microbial Endophyte of Crop Plant and Their Role in Plant Growth Promotion. Ph.D Thesis. University of Agricultural Science Bangalore. Bangalore. pp. 266.
- Arx J.A. von, Guarro J. et Figueras M.J. (1986). The ascomycete genus *Chaetomium*. 162 pp. Berlin et Stuttgart.
- Barnett H.L. and B. B. Hunter (nd). “Illustrated Genera of Imperfect Fungi”, 2nd Edn., Burgess Publishing, Minneapolis, MN, 1956, pp. 218.
- Bayman P., et al. (1997). Variation in endophytic fungi from roots and leaves of *Lepanthes* (Orchidaceae). *New Phytol.* 135: 143 – 149.
- Blumenstein, K. (2010). *Characterization of endophytic fungi in the genus Ulmus: putative agents for the biocontrol of Dutch Elm Disease (DED)*. Diploma thesis, University of Kassel.
- Bougoure J.J. and J.D.W. Dearnaley (2005). The Fungal endophyte of dipodium Variegatum (Orchidaceae). *Australasin Mycologist.* 24(1): 15-19.
- Chen J. et al. (2013). Diversity and Taxonomy of Endophytic Xylariaceous Fungi from Medicinal Plants of Dendrobium (Orchidaceae). *PLoS ONE* 8(3):1-11.
- Clemente M. (2009). Orchid conservation and trade: are these concepts incompatible?. In: Pridgeon A M, Suarez JP (eds) *Proceedings of the Second Scientific Conference on Andean Orchids*. Universidad Técnica Particular de Loja, Loja, Ecuador, pp 46 -55.
- Crous, P. W., Verkley, G. J. M., Groenewald, J. Z. and Samson, R. A. (2009). *Fungal Biodiversity*. CBS Laboratory manual series. CBS-KNAW Fungal Biodiversity Center.
- Dandu Anitha et al. (2013). Isolation and Characterization of Endophytic Fungi from Endemic Medicinal Plants of Tirumala Hills. *International Journal of Life Science Biotechnology and Pharma Research.* 2(3): 367-373.
- Dai C.C., Yu B.Y., Xu Z.L., Yuan S. (2008). Screening of endophytic fungi that promote the growth of *Euphorbia pekinensis*. *African Journal of Biotechnology* . 7 (19):3505-3510.
- De Hoog. G.S.J. Guarro. J. Gene and M.J. Figueras. (2000). *Atlas of clinical fungi*. 2nd ed. Vol. 1. Centraalbureauvoorschimmel cultures. Utrecht. The Netherlands.
- Dreyfuss M, Chapela IH (1994). Potential of fungi in the discovery of novel, low-molecular weight pharmaceuticals. In: *The discovery of natural products with therapeutic potential*. (Ed.): V. P. Gullo. Butterworth- Heinemann, London, United Kingdom. pp. 49-80.
- Feldman, T.S., O'Brien, H. and Arnold, A.E. (2008) Moth dispersal of mycoparasites and endophytes associated with *Claviceps paspali* and the grass *Paspalum* (Poaceae). *Microbial Ecology* 56: 742-750.
- Gange A.C.S., Dey A.F., Currie, B.V.C. Sutton (2007). Site and species differences in endophyte occurrence in two herbaceous plants. *J. of Ecol.* 95(4): 614-622.
- Gao Y. et al. (2011). Characterization of Five Fungal Endophytes Producing Cajaninstilbene Acid Isolated from Pigeon Pea [*Cajanus cajan* (L.) Millsp.]. *PLoS ONE* 6(11):1-9.
- Gezgin Y. and R. Eltem. (2009). Diversity of endophytic fungi from various Aegean and Mediterranean orchids (saleps). *Tubitak.* 33(2009):439-445.

- Gimenez, C., Cabrera, R., Reina, M., and Gonzalel-Coloma, A. (2007). Fungal endophytes and their role in plant protection. *Current Organic Chemistry* 11:707-720.
- Glenn A, Bodri M.S. (2012) Fungal Endophyte Diversity in Sarracenia. *PLoS ONE*. 7(3):1-7.
- Hamayun M. et al. (2010). Gibberellin production and plant growth promotion from pure cultures of *Cladosporium* sp. MH-6 isolated from cucumber (*Cucumis sativus* L.). *Mycologia*, 102(5): 989–995.
- Hamilton CE, Bauerle TL (2012). A new currency for mutualism? Fungal endophytes alter antioxidant activity in hosts responding to drought. *Fungal Divers* 54: 39–49.
- Hanada R.E. et al. (2010). Endophytic fungal diversity in *Theobroma cacao* (cacao) and *T. grandiflorum* (cupuac, u) trees and their potential for growth promotion and biocontrol of black-pod disease. *Fungal Biology*. 114 (2010): 901-910.
- Helander, M., Ahlholm, J., Sieber, T. N., Hinneri, S. and Saikkonen, K. (2007). *Fragmented environment affects birch leaf endophytes*. *New Phytologist* 175: 547- 553.
- Herre A.D., L.C. Meji' A, D.A. Kylo, E. Rojas, Z. Maynard, A. Bulter AND Sunshine A. V. (2007). Ecological Implication of Anti-Pathogen Effect of Tropical Fungal Endophytes and Mycorrhizae.. *Ecology*, 88(3): 550–558.
- Higgins, K.L., Arnold, A.E., Miadlikowska, J., Sarvate, S.D. and Lutzoni, F. (2007). Phylogenetic relationships, host affinity and geographic structure of boreal and arctic endophytes from three major plant lineages. *Mol. Phyl. and Evol.* 42(2): 543-555.
- Jantawong J., N. Potapohn, and O. Chatsrirung (2010). Seasonal Effect on Endophytic Fungal Diversity in *Geodurum* root. *Agriculture Journal*. 21(1): 35-42.
- Jeremy Martin O. Torres and Thomas Edison E. dela Cruz. http://biowikifarm.net/studienstiftung/Talk:Main_Page . 13 January 2014.
- Kasmir J., S.R.Senthilkumar, S.John Britto, and L. Joelri Michael Raj (2011). Identification of fungal endophytes from Orchidaceae members based on nrITS (Internal Transcribed Spacer) region. *International Research Journal of Biotechnology*. 2(6) 139-144.
- Khan S.A. et al. (2012). Isolation of Plant Growth Promoting Endophytic Fungi from Dicots Inhabiting Coastal Sand Dunes of Korea. *Pak. J. Bot.*, 44(4): 1453-1460.
- Kogel, K.-H., P. Franken, and R. H. (2006). Kelhoven, Endophyte or parasite - what decides?[J]. *Current Opinion in Plant Biology*, 9(4):358-363.
- Kuldau, G. and Bacon , C. (2008). Clavicipitaceous endophytes: Their ability to enhance resistance of grasses to multiple stresses. *Biological Control* 46(1): 57-71.
- Kumar A, Patil D, Rajamohan PR, Ahmad A (2013). Isolation, Purification and Characterization of Vinblastine and Vincristine from Endophytic Fungus *Fusarium oxysporum* Isolated from *Catharanthus roseus*. *PLoS ONE* 8(9):1-10.
- Kumar, D.S.S. and Hyde, K.D. (2004). Biodiversity and tissue-recurrence of endophytic fungi in *Tripterygium wilfordii*. *Fungal Diversity* 17: 69-90.
- Li, J.H. et al. (2008). Genetic diversity and potential for promotion of plant growth detected in nodule endophytic bacteria of soybean grown in Heilongjiang province of China[J]. *Soil Biology and Biochemistry*, 40(1):238-246.
- Lopez-Llorca, L.V., Jansson, H-B., Vicente, J.G.M. and Salinas, J. (2006). *Nematophagous* fungi as root endophytes. In Schulz B, Boyle C, Sieber T, eds. *Microbial root endophytes*. Berlin, Germany: Springer-Verlag. Pp.191-206.
- Mangunwardoyo W. and Suciati Mih, Gandjar I. (2012). Frequency of endophytic fungi isolated from *Dendrobium crumenatum* (Pigeon orchid) and antimicrobial activity. *Biodiversitas* 13: 34-39.
- McGinnis, M.R. (1980). *Laboratory handbook of medical mycology*. Academic Press, London, UK.
- Melfei E. Bungihan, Maribel G. Nonato, Siegfried Draeger, Scott Franzblau and Thomas Edison E. dela Cruz (2013). Antimicrobial and Antioxidant Activities of Fungal Leaf Endophytes Associated with *Pandanus amaryllifolius* Roxb. *Philippine Science Letters*. 6(2): 128-137.

- Min-Yuan Ho, Wen-Chuan Chung, Hung-Chang Huang , Wen-Hsia Chung and Wen-Hsin Chung (2012). Identification of Endophytic Fungi of Medicinal Herbs of *Lauraceae* and *Rutaceae* with Antimicrobial Property. *Taiwania*, 57(3): 229-241.
- Waqas M., Abdul Latif Khan, Muhammad Kamran, Muhammad Hamayun, Sang-Mo Kang, Yoon-Ha Kim and In-Jung Lee (2012). Endophytic Fungi Produce Gibberellins and Indoleacetic Acid and Promotes Host-Plant Growth during Stress. *Molecules* 2012(17) 10754-10773.
- Ovando I. et al. (2005). Isolation of Endophytic Fungi and Their Mycorrhizal Potential for the Tropical Epiphytic Orchids *Cattleya skinneri*, *C. aurantiaca* and *Brasavola nodosa*. *Asian Journal of Plant Science*. 4(3): 309-315.
- Paszkowski, U. (2006) Mutualism and parasitism: the yin and yang of plant symbioses. *Curr. Opin. Plant Biol.* 9(4): 364-370.
- Paul N. C., Jian X. D., Hyun K.S., Young P.C. and Seung H.Y. (2011). Distribution and Antifungal Activity of Endophytic Fungi in Different Growth Stages of Chili Pepper (*Capsicum annuum* L.) in Korea. *The Plant Pathology Journal*. 28(1) : 10-19.
- Perkins, A.J., Masuhara, G. & McGee, P.A. (1995). Specificity of the associations between *Microtis parviflora* (Orchidaceae) and its mycorrhizal fungi. *Australian Journal of Botany*: 43: 85-91.
- Purahong W, Hyde KD (2011). Effects of fungal endophytes on grass and nongrass litter decomposition rates. *Fungal Divers* 47: 1–7.
- Qadri M., S. Johri1, B. A. Shah, A. Khajuria, T. Sidiq, S. K. Lattoo, M. Z Abdin and S. Riyaz-Ul-Hassan (2013). Identification and bioactive potential of endophytic fungi isolated from selected plants of the Western Himalayas. *SpringerPlus*. 2(8): 1-14.
- R.A. Cating, A. J. Palmateer, C. M. Stiles, P. A. Rayside, and D. A. Davison (2013). Black Rot of Orchids Caused by *Phytophthora palmivora* and *Phytophthora cactorum*. Source: <http://edis.ifas.ufl.edu>.
- Rodriguez, R., White, J., Arnold, A. and Redman, R. (2009) Fungal endophytes: diversity and functional roles. *New Phytologist* 182(2): 314-330.
- Rola S. Mahmoud, Kazuhiko Narisawa (2013). A New Fungal Endophyte, *Scolecobasidium humicola*, Promotes Tomato Growth under Organic Nitrogen Conditions. *Plos One*. 8(11):1-8.
- Rudgers, J.A., Afkhami, M.E., Rua, M.A., Davitt, A.J., Hammer, S., and Huguet, V.M. (2009). A fungus among us: broad patterns of endophyte distribution in the grasses. *Ecology* 90:1531- 1539.
- Saitong P., Panthawee W., Stonsaovapak S., and Li C. (2010). Isolation and primary identification of endophytic fungi from *Cephalotaxus mannii* trees. *Maejo International Journal of Science and Technology*. Thailand, 4(03): 446-453.
- Sawmya K., Vasudevan T.G., Murali T.S. (2013). Fungal endophytes from two orchid species – pointer towards organ specificity. – *Czech Mycol.* 65(1): 89–101.
- Schulz, B. and Boyle, C. (2005). The endophytic continuum. *Mycological Research* 109: 661-686.
- Schulz, B. and Boyle, C. (2006). What are endophytes? In: Schulz, B.J.E., BOYLE, C.J.C. AND SIEBER, T.N., (eds.) *Microbial Root Endophytes*, Springer-Verlag, Berlin.,pp.1–13.
- Sibounnavong, P., Sysouphan, P., Xay Ly, Phoutsay, P., Soyong, K., Promrin, K., Pongnak, W., and K. Soyong. (2006). Application of biological products for organic crop production of kangkong (*Ipomoea aquatica*). *An International Journal of Agricultural Technology* 2(2):177-189.
- Stone, J. K., J. F. White, Jr., J. D. Polishook (2004). Endophytic fungi. In: *Measuring and Monitoring Biodiversity of Fungi. Inventory and monitoring methods*. G. Mueller, G. Bills, M. Foster (eds.). Elsevier Academic Press, Boston, MA. pp. 241-270.
- Strobel, G. and Daisy, B. (2003). *Microbiol.Mol. Biol. Rev.* 67: 491–502.

- Su Z-Z et al. (2013). Evidence for Biotrophic Lifestyle and Biocontrol Potential of Dark Septate Endophyte *Harpophora oryzae* to Rice Blast Disease. *PLoS ONE* 8(4): 1-14.
- Sun Y, Wang Q, Lu XD, Okane I, Kakishima M. (2011). Endophytic fungi associated with two *Suaeda* species growing in alkaline soil in China. *Mycosphere* 2(3): 239-248.
- Supaphon P., Phongpaichit S., Rukachaisirikul V. and Sakayaroj J. (2013). Antimicrobial Potential of Endophytic Fungi Derived from Three Seagrass Species: *Cymodocea serrulata*, *Halophila ovalis* and *Thalassia hemprichii*. *PLoS ONE* 8(8): 1-9.
- Sutthiwari P. (2006). Orchids Production. Rajamangala University Technology Tawan-ok Chantaburi campus. pp.111.
- Susheel Kumar and Nutan Kaushik (2013). Endophytic Fungi Isolated from Oil-Seed Crop *Jatropha curcas* Produces Oil and Exhibit Antifungal Activity. *plosone*. 8(2): 1-8.
- Susheel Kumar, Nutan Kaushik and Peter Proksch (2013). Identification of antifungal principle in the solvent extract of an endophytic fungus *Chaetomium globosum* from *Withania somnifera*. *SpringerPlus*. 2(37): 1-10.
- Tanaka, A., et al. (2006). Reactive Oxygen Species Play a Role in Regulating a Fungus-Perennial Ryegrass Mutualistic Interaction[J]. *Plant Cell*. 18(4): 1052-1066.
- Tao, G., Liu, Z.Y., Hyde, K.D. Lui, X.Z. and Yu, Z.N. (2008). Whole rDNA analysis reveals novel and endophytic fungi in *Bletilla ochracea* (*Orchidaceae*). *Fungal Diversity*. 33: 101-122.
- Tempesta S., A. Robini, F. Pupulin and A. Ramballi (2011). Pestalotiopsis endophytes from leaves of two orchid species collected in Costa Rica. *Cryotogamie mycologie*. 32(3): 315-321.
- Thamasiri K. (2005). Orchids Production Technology. Amarin Printing and Publication. Bangkok. pp. 283.
- Tiamphet N. and W. Nuangmek (2013). Screening of endophytic fungi for control *rhizoctonia solani* rot of eucalyptus seedling. *KHON KAEN AGR. J.* 1(2013): 505-512.
- Valdes M., Guerrero H.B., Martinez L. and Rafael H. Viquez (2011). The root colonizing fungi of the terrestrial orchid *Cypripedium irapeanum*. *LANKESTERIANA* 11(1): 15-21.
- Vega, F.E. (2008). Insect pathology and fungal endophytes. *Journal of Invertebrate Pathology*. 98: 277-279.
- Vega F. E., Simpkins, A., Aime, M.C., Posada, F., Peterson, S.W., Stephen A., Rehner, S.A., Infante, F., Castillo, A. and Arnold, A. E. (2010). Fungal endophyte diversity in coffee plants from Colombia, Hawai'i, Mexico and Puerto Rico. *Fungal Ecology*. 3: 122-138 .
- Yaun, Z., Zhang, C., and Lin, F. (2009). Role of diverse non-systemic fungal endophytes in plant performance and response to stress: progress and approaches. *Journal of Plant Growth Regulation* 29:116-126.
- Yin Lu, Chuan Chen, Hong Chen, Jianfen Zhang and Weiqin Chen (2012). Isolation and Identification of Endophytic Fungi from *Actinidia macrosperma* and Investigation of Their Bioactivities. *Hindawi Publishing Corporation*:1-8.



This material is reserved for educational use only, not allowed for commercial use.

Forbidden to modify the content, and cite the document when use. **ศรายุทธ ผลโพธิ์ วรรณะ ต้ว**

Sour V., Phonpho S and Soyong K. (2015) Isolation of endophytic fungi from some orchid varieties. *Journal of Agricultural Technology* 11(5):1243-1254



Isolation of endophytic fungi from some orchid varieties

Sour V.^{*}, Phonpho S. and Soyong K.

Faculty of Agricultural Technoloty, King Mongkut's Institute of Technology Ladkrabang, Bangkok, Thailand

Sour V., Phonpho S and Soyong K. (2015) Isolation of endophytic fungi from some orchid varieties. Journal of Agricultural Technology 11(5):1243-1254

Endophytic fungi from orchids in Chantaburi province was studied. Endophytic fungi were isolated from 9 orchid varieties in greenhouse condition. Leaves and roots of *Grammatophyllum scriptum*, *Cymbidium dayanum*, *Dendrobium hercoglossum*, *Dendrobium palpebrae*, *Torenia journeri*, *Doritis pulcherrima*, *Dendrobium crumenatum*, *Dendrobium friedericksianum* and *Grammatophyllum specinocum* were collected. All samples were collected then brought to laboratory. Isolation was done by surface sterilization, removed the outer surface and got inner tissues then place on water agar and transferred culture to potato dextrose agar to be pure culture. The morphological identification based on colony and agar color, liquid drops, colony shapes, the growth rate and spores characteristic Twelve species are identified as *Chaetomium cochliodes*, *Chaetomium cupreum*, *Xylaria* sp., *Colletotrichum* sp., *Nigrospora* sp., *Phoma* sp. *Curvularia* sp., *Fusarium* sp., *Pestalotiopsis* sp., *Corynascus* sp. and *Cladosporium*-like fungi.

Keywords: Endophytic fungi, Orchids

Introduction

Plants are always colonized with other microorganisms. Endophytic fungi are one kind of microorganism that live within healthy plant tissue causing no harm to the host plant. The meaning of endophytic fungi is so broad. Modern usage of the term endophytic fungi in mycology refers to those fungi which live within leaves stems roots and other part of apparently healthy host plants and can not be seen in visually signs of infection (Stone J.K. *et al.*, 2004). Dreyfuss and Chapela (1994) estimated that there may be at least one million species of endophytic fungi alone. It means that almost all plant species are usually infected with endophytic fungi.

^{*} Coressponding Author: E-mail address: sourvannak@hotmail.com

Nowaday endophytic fungi take attention from many scientists as estimated that such species may useful as sources of anticancer, antidiabetic, insecticidal and immunosuppressive compounds (Strobel, G. and Daisy, B., 2003). Endophytic fungi from medicinal plants can be used for the development of drugs. For example *Fusarium oxysporum* isolated from *Catharanthus roseus* plant can produce Vinblastine and Vincristine, excellent anti-cancer drugs (Kumar A. *et al.*, 2013).

Orchids are monocotyledonous plants and taxonomically belong to Orchidales and Orchidaceae. There are 15000 to 25000 of orchid species distributed all around the world, mostly can found in tropical environment. (Thamasiri K., 2005). Fungal endophytes have now been investigated in a large number of orchid species from around the world (Perkins *et al.* 1995). The fungi involved with orchids are almost all members of the phylum Basidiomycota group. The results may vary with the species of orchid.

Materials and methods

Plant sample collection

Roots and leaves of 9 orchid varieties (*Grammatophyllum scriptum*, *Cymbidium dayanum*, *Dendrobium hercoglossum*, *Dendrobium palpebrae*, *Torenia fournieri*, *Doritis pulcherrima*, *Dendrobium crumenatum*, *Dendrobium friedricksianum* and *Grammatophyllum specinocum*). Sample were collected in greenhouse from Rajamangala University Technology Tawan-ok Chantaburi Campus, Chantaburi province, Thailand on 1st December 2014.

All samples were placed in sterile polythene bags and brought to laboratory in an icebox and stored at 4.5 °C before isolating endophytic fungi within 48 hours.

Isolation and identification of endophytic fungi

The surface sterilization was done by procedure used by Blumenstein (2010) with some modifications. Symptomless roots of each plant species were cut into 2–3 mm sections were rinsed in a sequence distilled water for 1 min, 75% ethanol for 30s, 3% NaClO for 1 min, and 75% ethanol for 30s and then rinsed in sterile distilled water three times. Each root was placed on water agar (WA). The isolates were subcultured on potato dextrose agar for morphological identification and conidial production.

The endophytic fungi were identified initially according to their colony, mycelium and spore characteristics according to the discription of Barnett and Hunter (1998) and Von Arx (1978).

Results and Discussion

Many isolates can't identify the species because it doesn't produce spores. Twelve species belonging to Ascomycota were isolated from 9 orchid varieties. They are as follow:

1. *Chaetomium cochliodes*

Habitat: *Grammatophyllum scriptum*, *Cymbidium dayanum*, *Dendrobium hercoglossum* and *Torenia fournieri*.

The colonies are quit slowly growing, for cultured on PDA medium in room temperature for 7 days have 7 cm in diameter, cottony and white in color initially and then become soft-green. *C. cochliodes* has distinctive small brown 'lemon' or 'football'shaped ascospores. (Arx, J.A. von *et al.* 1986).

This species also isolated as endophytic fungi from *Cirsium arvense*, (Gange AC. Et al., 2012). Li Gy et al. 2006 indicated that epipolythiodioxopiperazines and chaetocochin were isolated from the ethyl acetate extract of the solid-state fermented rice culture of the fungus *Chaetomium cochliodes*. Their structures were elucidated on the basis of spectroscopic analysis. Compounds 1, 3, and 4 exhibited significant cytotoxicity in vitro against cancer cell lines Bre-04, Lu-04, and N-04,




Fig. 1: *Chaetomium cochliodes* A: The colony that culture on PDA 30 days, B The spores under microscops 10X C The ascospores under microscops 40X

2. *Chaetomium cupreum*

Habitat: *Grammatophyllum scriptum*.

The growth was quite slow, cultured on PDA medium for 7 days the growth have 4.4 cm in diameter. The hyphae are red and it produces the red color around mycelia.

Septate hyphae, perithecia, asci and ascospores are visualized. Perithecia are large, red color, fragile, globose to flask shaped and have filamentous, hair-like, brown to black appendages (setae) on their surface. Perithecia have ostioles (small rounded openings) and contain asci and ascospores inside (De Hoog *et al.*, 2000).

This species was isolated as endophytic fungus from *Macleaya cordata* and have ability to antifungal activity against *Rhizoctonia solani*, *Botrytis cinerea*, *Pytium ultimum*, and antitumor activities against HL-60 and A549. It is toxic to *Artemia salina* (Bi Z.M. *et al.*, 2010).




Fig. 2: *Chaetomium cupreum* A: The colony cultured on PDA 30 days B: The spores under microscopes 10X and C: The ascospores under microscopes 40X

3. *Xylaria* sp.

Habitat: *Cymbidium dayanum*, *Dendrobium hercoglossum*, *Doritis pulcherrima*, *Dendrobium crumenatum*, *Dendrobium friedericksianum* and *Grammatophyllum specinocum*.

The fungi have white mycelia; the growth was quite slow, for cultured on PDA in room temperature for 7 days have 5.8 cm in diameter. They produce the stroma on 20th day but didn't produce ascus and ascospore.

Xylaria sp. was the common species that can isolated from other orchid varieties. Previous research also found that this species are most colonized in 7 *Dendrobium* sp. in China (Chen J. *et al.* 2013). Sawmya K. *et al.* (2013) also isolated *Xylaria* sp. from *Bulbophyllum neilgherrense* and *Pholidota pallida*.




Fig. 3: *Xylaria* sp. A: Colony culture on PDA 30 days B: Mycelia of xylaria sp. under microscopes 40X

4. *Colletotrichum* sp.

Habitats: *Cymbidium dayanum*, *Dendrobium hercoglossum*, *Torenia fournieri*, *Dendrobium crumenatum* and *Grammatophyllum specinocum*.

The growth on PDA medium 7 days have 8.5-9 cm, the mycelia initially white-grey and then become black-brown. The conidia have no color with rods shape (Busarakham K.,2001).

Previous study from India showed that the endophytic fungus, *Colletotrichum gloeosporiodes* was isolated from *Justicia gendarussa*, a medicinal plant have ability to produce Taxol that is a potent anticancer drug used widely in the treatment of a variety of cancers (Gangadevi V. and Muthumary J., 2008).




Fig. 4: *Colletotrichum* sp. A: The colony culture on PDA 30 days and B: The spores under microscope 40X

5. *Nigrospora* sp.

Habitat: *Dendrobium hercoglossum*

Nigrospora grows quit fast and produces woolly colonies on potato dextrose agar (PDA) at room temperature, on 7th days has 9 cm in diameter. Color of the colony is white initially and then becomes gray with black areas and turns to black eventually from both front and reverse. Spores (conidia) are produced singly on swollen urn-shaped conidiophores and are egg-shaped to

flattened-spherical, black, and often have an equatorial colourless line or germ slit.

This species also can isolated from deep sea environment at Arabian Sea and The secondary metabolites produced from this organism showed potent antimicrobial and anticancer activities with immediate application to cosmetics and pharmaceutical industries. (Arumugam GK, *et al*, 2015)




Fig. 5: *Nigrospora* sp. **A:** The colony culture on PDA 30 days **B:** The hyphea and spores under microscope 40X

6. *Phoma* sp.

Habitats: *Cymbidium dayanum*, *Doritis pulcherrima* and *Grammatophyllum specinocum*.

The hyphae were growing rapidly, for 2 days that cultured on PDA medium have 6.5 cm in diameter. The hyphae first growth have white color and than become black-brown. Spores are colorless and unicellular. The pycnidia are black and depressed in the tissues of the host.

Phoma sp. are endophytic fungi that can isolated from many plants. Wang LW *et al.* (2012) isolated *phoma* sp. from chinese medicinal plant *Arisaema erubescens* and showed that the bioactive metabolites from this species can inhibited the growth of four plant pathogenic fungi (*Fusarium oxysporium*, *Rhizoctonia solani*, *Colletotrichum gloeosporioides*, and *Magnaporthe oryzae*). This species also isolated from roots of cucumber in Korea and showed that have ability to produce Gibberellins and Indoleacetic Acid and Promotes Host-Plant Growth during Stress (Waqas M. *et al.*, 2012).




Fig. 6: *Phoma* sp. A: The colony culture on PDA 30 days and B: The spores and mycelia under microscope 40X

7. *Curvularia* sp.

Habitats: *Grammatophyllum scriptum* and *Doritis pulcherrima*.

The growth was quit fast for cultured on PDA medium for 2 days have 5.2cm in diameter. From the front, the color of the colony is white to pinkish gray initially and turns to black as the colony matures. From the reverse, it is dark brown to black. Conidia are pale brown, with three or more transverse septa (phragmoconidia) and are formed apically through a pore (poroconidia) in a sympodially elongating geniculate conidiophore similar to *Drechslera*. Conidia are cylindrical or slightly curved, with one of the central cells being larger and darker. Germination is bipolar and some species may have a prominent hilum (McGinnis, M.R. 1980).

Zakaria L. *et al.* (2010) isolated this species from *Oryza sativa*. Aharwal R.P. *et al.* (2014) also isolated this fungus from *Calotropis procera* (Linn.) R.Br. a widely used medicinal plant in India.




Fig. 7: *Curvularia* sp. A: The colony culture on PDA 30 days and B: The spores under microscope 40X

8. *Fusarium* sp.

Habitat: *Grammatophyllum scriptum*, *Cymbidium dayanum*, *Dendrobium palpebrae*, *Torenia fournieri* and *Dendrobium friedericksianum*.

The hyphae are white. The growth was quite fast, for cultured on PDA in room temperature, for 4 days have 4.9 cm in diameter. The conidia are fusiform, slightly curved, pointed at the tip. *Fusarium* is a filamentous fungus widely distributed on plants and in the soil. It is found in normal mycoflora of commodities, such as rice, bean, soybean, and other crops.

Previous study showed that *Fusarium redolens*, isolated from Himalayan yew can produce taxol, the anticancer drug (Garyali S, 2013). Sogra F.M. and Raj M.B.(2014) isolated *Fusarium oxysporum* from leaf of *Nothapodytes foetida*, ethyl acetate extract from this fungus show ability to inhibited the growth of *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Candida albicans*.




Fig. 8: *Fusarium* sp. A: The colony culture on PDA 30 days and B: The spores under microscope 40X

9. *Pestalotiopsis* sp.

Habitat: *Dendrobium friedericksianum*, *Cymbidium dayanum* and *Grammatophyllum scriptum*.

The mycelia are white in color. This species have quite slow of growing. For growing on PDA in room temperature for 7 days have 5.5cm in diameter. Conidiophores (annellides) produced within compact fruiting structures (aecervuli or pycnidia). Spores (conidia) 4 to 5 celled, with the two or three central cells dark brown, and with two or more apical appendages or hairs.

Pestalotiopsis sp. is the common endophytic fungi that can isolate from orchids and other plants. They also isolated from other orchid varieties from other country such as Tempesta S. *et al.* (2011) isolated endophytic fungi from leaves of *C. mrcranthum* and *O. adendrobium* also found 29 strains of *Pestalotiopsis* sp. Wei J.G *et al.* (2007) isolated this species that associated with plants of Podocarpaceae, Theaceae and Taxaceae in southern China also obtained 24 species. Kumaran RS *et al.* (2010) isolated *Pestalotiopsis ver-*

sicolor and *Pestalotiopsis neglecta* as endophytic fungi from Japanese Yew tree, *Taxus cuspidata* and indicated that they can produce taxol concentration induces increased cell death in cancer.




Fig. 9: *Pestalotiopsis* sp. A: The colony culture on PDA 30 days B: The conidia under microscopes 40X

10. *Corynascus* sp.

Habitat: *Cymbidium dayanum*

The fungus have quit fast in growth. Only 3 days the mycelia have 9.0 cm in diameter. It has black-white mycelia, produce lots of black spores and inside spores there are lots of little ascospores.




Fig. 10: *Corynascus* sp. A: The colony culture on PDA 30 days and B: The spores and ascospores under microscopes 40X

11. *Cladosporium*-like fungi

Habitat: *Grammatophyllum scriptum*, *Cymbidium dayanum*, *Dendrobium palpebrae*, *Doritis pulcherrima* and *Dendrobium crumenatum*

The initial growth the mycelia have white color and then become black. The growth was quit fast. On 6th day of culture on PDA have 9 cm in diameter. The mycelia have branches and the lots of small spore.




Fig. 11: *Cladosporium*-like fungi A: The colony culture on PDA 30 days and B: The spores and mycelia under microscopes 40X

12. *Achaetomium* sp.

Habitat: *Grammatophyllum scriptum*

The mycelia growth look like the yellow sheets and this fungus growth quit slow, cultured on PDA medium in room temperature for 7 days have 5.5 cm in diameter. The spores are large with yellow in color. Inside the spores have many ascospores.



Fig. 12: *Achaetomium* sp. A: The colony culture on PDA 30 days B: The spores under microscopes 10X and C: The ascospores under microscopes 40X

Acknowledgement

This work is the part of my thesis and has got scholarship from King Mongkut's Institute of Technology Ladkrabang. I would like to express my sincere to thank to Faculty of Agriculture Technology, King Mongkut's Institute of Technology Ladkrabang (KMITL), Bangkok, Thailand. Thank to my advisor Dr. Sarayut Phonpho, my co-advisor Dr. Kasem Soytung and other teachers and friends who help me to do this research.

References

- Aharwal R.P., Kumar S. and S.S. Sandhu. (2014). Isolation and antibacterial property of endophytic fungi isolated from Indian dedicial plant *Calotropis procera* (Linn.) R.Br. World journal of phammacy and pharmaceutical sciences. 3(5): 678-691.

- Arumugam G.K., Srinivasan S.K., Joshi G., Gopal D., Ramalingam K. (2015). Production and characterization of bioactive metabolites from piezotolerant deep sea fungus *Nigrospora* sp. in submerged fermentation. *J Appl Microbiol.* 118(1): 99-111.
- Arx J.A. von, Guarro J. et Figueras M.J. (1986). The ascomycete genus *Chaetomium*. 162 pp. Berlin et Stuttgart.
- Barnett H.L. and B. B. Hunter (nd). "Illustrated Genera of Imperfect Fungi", 2nd Edn. Burgess Publishing, Minneapolis, MN, 1956, pp. 218.
- Bi Z.M., Chao H., Yang G.M., Chen Y.Z. and Chen S.Y. (2010). Separation and determination of the bioactivity of oosporein and determination of the bioactivity of oosporein from *Chaetomium cupreum*. *African Journal of Biotechnology.* 9(36): 5955-5961.
- Blumenstein, K. (2010). Characterization of endophytic fungi in the genus *Ulmus*: putative agents for the biocontrol of Dutch Elm Disease (DED). Diploma thesis. University of Kassel.
- Busarakham K. (2001). Diseases of Terrestrial Orchids, Endophytic Fungi on Leaves and Roots and Diversity and Rhizosphere soil. Master's degree thesis. Kasetsart University. Bangkok, Thailand. pp 259.
- Chen J., Zhang L.C., Xing Y.M., Wang Y.Q., Xing X.K., Zhang D.W., Liang H.Q. and Guo S.X. (2013). Diversity and taxonomy of endophytic xylariaceous fungi from medical plants of *Dendrobium* (Orchidaceae). *Plos one.* 8(3): 1-11.
- De Hoog. G.S.J. Guarro. J. Gene and M.J. Figueras. (2000). Atlas of clinical fungi. 2nd ed. Vol. 1. Centraalbureau voor schimmel cultures. Utrecht. The Netherlands.
- Dreyfuss M, Chapela IH. (1994). Potential of fungi in the discovery of novel, low-molecular weight pharmaceuticals. In: The discovery of natural products with therapeutic potential. (Ed.): V.P. Gullo. Butterworth-Heinemann. London. United Kingdom. pp.49-80.
- Gangadevi V. and Muthumary J. (2008). Isolation of *Colletotrichum gloeosporioides*, a novel endophytic fungi taxol-producing fungus from the leaves of a medicinal plant, *Justicia gendarussa*. *Mycologia Balcanica.* 5:1-4.
- Gange A.C., Eschen R., Weam J.A., Thawer A. and Sutton B.C. (2012). Differential effects of folia endophytic fungi on insect herbivores attacking a herbaceous plant. *Oecologia.* 168(4): 1023-1031.
- Garyali S., Kumar A., and Reddy MS. (2013). Taxol production by an edophytic fungus, *Fusarium redolens*, isolated from Himalayan yew. *J Microbiol Biotechnol* 23(10): 1372-80.
- Kumar A., Patil D., Rajamohanan PR and Ahmad A. (2013). Isolation, purification and characterization of vinblastine and Vincristine from endophytic fungus, *Fusarium oxysporum*, isolated from *Catharanthus roseus*. *Plos one.* 8(9): 1-10.
- Kumaran RS. Kim H.J. and Hur B.K. (2010). Taxol-producing [corrected] fungal edophyte, *Pestalotiopsis* species isolated from *Taxus cuspidate*. *Biosci Bioeng.* 110(5): 541-6.
- Li G.Y., Li B.G., Yang T., Yan J.F., Liu G.Y. and Zhang G.L. (2006). Chaetocochins A-C, epipolythiodioxopiperazines from *Chaetomium cochliodes*. *J Nat Prod.* 69(9): 1374-6.
- McGinnis, M.R. (1980). Laboratory handbook of medical mycology. Academic Press, London, UK.
- Perkins A.J., Masuhara G. and McGee P.A. (1995). Specificity of the associations between *Microtis parviflora* (Orchidaceae) and its mycorrhizal fungi. *Australian Journal of Botany.* 43: 85-91.
- Rodriguez R. and Redman R. (2008). More than 400 million years of evolution and some plants still can't make it on their own: plant stress tolerance via fungal symbiosis. *Journal of experimental botany.* 59(5): 1109-4.

- Sawmya K., Vasudevan T.G. and Murali T.S. (2013). Fungal endophytes from two orchid species-pointer towards organ specificity. *Czech mycol.* 65(1): 89-101.
- Sogra F.M. and Raj M.B. (2014). A study on the antimicrobial potentials of an endophytic fungus *Fusarium oxysporum* NFX 06. *Journal of medical and bioengineering.* 3(3): 162-166.
- Stone J.K., White J.F., Polishook J.D. (2004). Endophytic fungi. In: *Measuring and monitoring biodiversity of fungi. Inventory and monitoring methods.* G. Mueller, G. Bills, and M. Foster (eds.). Elsevier academic press, Boston, MA. Pp. 241-270.
- Strobel G. and Daisy B. (2013). *Microbiol Mol Biol.* 67: 491-502.
- Thamasiri K. (2005). *Orchids production technology.* Amarin printing and publication. Bangkok. pp. 283.
- Tempesta S., Robini A., Pupulin F., and Rambulli A. (2011). *Pestalotiopsis* endophytes from leaves of two orchid species collected in Costa Rica. *Crytogamie mycologie.* 32(3): 315-321.
- Gangadevi V. and Muthumary J. (2008). Isolation of *Colletotrichum gloeosporioides*, a novel endophytic taxol-producing fungus from the leaves of a medicinal plant, *Justicia gendarussa*. *Mycologia balcanica.* 5: 1-4.
- Von Arx J.A. (1978). The genera of fungi sporulation in pure culture. AR Gantner verlag KG (eds). FL-9490, Vaduz.
- Wang L.W., Xu B.G., Wang J.Y., Su Z.Z., Lin F.C., Zhang C.L. and Kubicek C.P. (2012). Bioactive metabolites from *Phoma* species, an endophytic fungus from the Chinese medicinal plant *Arisaema erubescens*. *Appl microbial biotechnol.* 93(3): 1231-9.
- Waqas M., Abdul L., Kamran M., Hamayun M., Kang S., Kim Y. and Lee L. (2012). Endophytic fungi produce gibberellins and indoleacetic acid and promotes host-plant growth during stress. *Molecules.* 17: 10754-10773.
- Wei J.G., Xu T., Guo L.D., Liu A.R., Zhang Y. and Pan X.H. (2007). Endophytic *Pestalotiopsis* species associated with plants of podocarpaceae, theaceae and taxaceae in southern China. *Fungal diversity.* 24: 55-74.
- Zakaria L., Yaakop A.S., Salleh B. and Zakaria M. (2010). Endophytic fungi from paddy. *Trop life sci res.* 21(1): 101-107.

(Received xxxxxx, accepted xxxxxx)

Isolation of Endophytic Fungi from some Orchid Varieties

ประวัติหัวหน้าโครงการ

ชื่อ-สกุล ชื่อ นาย ศรายุทธ ผลโพธิ์

เพศ ชาย หญิง วันเดือนปีเกิด

ตำแหน่งปัจจุบัน ผู้ช่วยศาสตราจารย์

หลักสูตรพืชสวน สาขาวิชาเทคโนโลยีการผลิตพืช คณะเทคโนโลยีพระจอมเกล้าเจ้าคุณทหาร

ลาดกระบัง โทรศัพท์ 02-3264318 โทรสาร 02-3264318 e-mail kpsarayu@kmitl.ac.th

ประวัติการศึกษา

ปีที่จบการศึกษา	ระดับปริญญาตรีโท เอก และประกาศนียบัตร	อักษรย่อปริญญาและชื่อเต็ม	สาขาวิชา	วิชาเอก	ชื่อสถาบันการศึกษา	ประเทศ
2534	ตรี	วท.บ (เกษตรศาสตร์) เกียรตินิยม	พืชสวน	พืชสวน	มหาวิทยาลัยเกษตรศาสตร์	ไทย
2539	โท	วท.ม (เกษตรศาสตร์)	พืชสวน	พืชสวน	มหาวิทยาลัยเกษตรศาสตร์	ไทย
2549	เอก	วท.ค (พืชสวน)	พืชสวน	พืชสวน	มหาวิทยาลัยเกษตรศาสตร์	ไทย

สาขาวิจัยที่มีความชำนาญพิเศษ Landscape design

ทุนการศึกษาและทุนวิจัยที่เคยได้รับ

ปี พ.ศ.	ทุนการศึกษาและทุนวิจัย	สถาบันที่ให้
2553	การสำรวจพรรณไม้ชายทะเลเพื่อใช้ในงาน	สถาบันเทคโนโลยีพระจอมเกล้าเจ้าคุณ
2557	ออกแบบภูมิทัศน์ ริมฝั่งทะเลอันดามันในบางพื้นที่ของประเทศไทย	ทหารลาดกระบัง
	การพัฒนาาระบบสวนแนวตั้งเพื่อประดับอาคาร	สถาบันเทคโนโลยีพระจอมเกล้าเจ้าคุณทหารลาดกระบัง

ผลงานวิจัย/งานสร้างสรรค์ที่ตีพิมพ์เผยแพร่

ศรายุทธ ผลโพธิ์และอลิศรา มีนะกนิษฐ. 2548. การประเมินศักยภาพแหล่งท่องเที่ยวทางเรือเพื่อวางแผนปรับปรุงสภาพภูมิทัศน์: เส้นทางแม่น้ำเจ้าพระยาดังแต่จังหวัดนนทบุรีถึงจังหวัดพระนครศรีอยุธยา. วารสารวิทยาศาสตร์เกษตร. ปีที่ 36 ฉบับที่ 5-6 (พิเศษ). หน้า 848-851.

ศรายุทธ ผลโพธิ์และอลิศรา มีนะกนิษฐ 2548. ลานโล่งในเมือง: กรณีศึกษาหน้าห้างสรรพสินค้าเดอะมอลล์บางกะปิ. วารสารวิทยาศาสตร์เกษตร. ปีที่ 36 ฉบับที่ 5-6 (พิเศษ). หน้า 856-859.

This material is reserved for educational use only, not allowed for commercial use.

Forbidden to modify the content, and cite the document when use. ศรายุทธ ผลโพธิ์ วรรณะ สัว

- ศรายุทธ ผลโพธิ์. 2549. การออกแบบปรับปรุงภูมิทัศน์โรงเรียนราชวินิตมัธยมเพื่อรวบรวมพันธุ์ไม้มงคล
ในโครงการอนุรักษ์พันธุกรรมพืชอันเนื่องมาจากพระราชดำริ สมเด็จพระเทพรัตนราชสุดาฯสยาม
บรมราชกุมารี. วารสารวิทยาศาสตร์เกษตร. ปีที่ 37 ฉบับที่ 6 (พิเศษ). หน้า 917-920.
- ศรายุทธ ผลโพธิ์ อติศรา มีนะกนิษฐ เอี่ยมพร วิสมหมาย และเอกรินทร์ อนุกุลยุทธชน. 2548. ความคิดเห็น
ของนักท่องเที่ยวดำรงเท้าเที่ยวทางน้ำเส้นทางแม่น้ำเจ้าพระยาระหว่างกรุงเทพมหานครถึงจังหวัด
พระนครศรีอยุธยา. วิทยาศาสตร์เกษตรศาสตร์ (สังคม). ปีที่ 27 ฉบับที่ 1. หน้า 67-78.
- เจษฎา แก้วฉาย นิจสิริ แววชาญ และศรายุทธ ผลโพธิ์. 2552. ความพึงพอใจต่อสภาพภูมิทัศน์และสิ่งอำนวยความสะดวก
ความสะดวกของผู้ใช้สถานที่ภายในพุทธมณฑลจังหวัดนครปฐม. วารสารเกษตรพระจอมเกล้า. ปี
ที่ 27 ฉบับที่ 2. หน้า 101-112.
- Sarayut Phonpho and Jessada Kaewchai. 2013. Landscape Planning and Design at BuddhamonthonPark,
Narathiwat, Thailand. Journal of Agricultural Technology. Vol. 9(2): 203-212.

